

**HUMAN CCL26 (68AA) PROTEIN****Cat.#:** 12022**Product Name:** Human CCL26 (68AA) Protein**Size:** 10 µg, 50 µg and 100 µg**Synonyms:** C-C Motif Chemokine 26;CC Chemokine IMAC;Eotaxin-3;Macrophage Inflammatory Protein 4-Alpha;MIP-4-Alpha;Small-Inducible Cytokine A26;Thymic Stroma Chemokine-1;TSC-1;CCL26;SCYA26**Target:** CCL26**UNIPROT ID:** Q9Y258**Description:** Recombinant Human C-C Motif Chemokine 26 is produced by our E.coli expression system and the target gene encoding Ser27-Leu94 is expressed.**Background:** Chemokine (C C Motif) Ligand 26 (CCL26) is a novel small cytokine belonging to the CC chemokine family, which involved in immunoregulatory and inflammatory processes. CCL26 is expressed constitutively in thymus, but only transiently in phytohemagglutinin-stimulated peripheral blood mononuclear cells. It specifically binds and induces chemotaxis in T cells and elicits its effects by interacting with the chemokine receptor CCR4. Eotaxin-3/CCL26, along with Eotaxin-1 and Eotaxin-2, selectively activates the CC chemokine receptor 3 (CCR3). The Eotaxin-3-CCR3 interaction may play an important role in allergic diseases such as atopic dermatitis and bronchial asthma. The full-length cDNA for Eotaxin-3 encodes a protein of 94 amino acids with a putative signal peptide of either 23 or 26 amino acid residues. Both the 71 and 68 amino acid residue variants of recombinant Eotaxin-3 demonstrate equal potency in inducing chemotaxis of a human CCR3-transfected cell line. Unlike most other CC chemokines, Eotaxin-3 maps to human chromosome 7q11.2, within 40 kilobases of the Eotaxin-2 loci. Eotaxin-3 and Eotaxin-2 are unique in that they are the only chemokines identified to date that map to chromosome 7.**Species/Host:** E.coli**Molecular Weight:** Predicted MW is 8.21 Kda. Protein runs at 13KDa under reducing conditions.**Molecular Characterization:** Not available**Purity:** Greater than 95% as determined by reducing SDS-PAGE.**Formulation & Reconstitution:** Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

**Storage & Shipping:** Store at  $-20^{\circ}\text{C}$  to  $-80^{\circ}\text{C}$  for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at  $-80^{\circ}\text{C}$  (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.




Figure 1. Greater than 95% as determined by reducing SDS-PAGE.