

HUMAN CD27 PROTEIN, MFC-HIS TAG

Cat.#: 11141

Product Name: Human CD27 Protein

Size: 10 µg, 50 µg and 100 µg

Synonyms: CD27;TNFRSF7;S152;T14;Tp55

Target: CD27

UNIPROT ID: P26842

Description: Recombinant human CD27 protein with C-terminal mouse Fc and 6xHis tag

Background: The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor is required for generation and long-term maintenance of T cell immunity. It binds to ligand CD70, and plays a key role in regulating B-cell activation and immunoglobulin synthesis. This receptor transduces signals that lead to the activation of NF-kappaB and MAPK8/JNK. Adaptor proteins TRAF2 and TRAF5 have been shown to mediate the signaling process of this receptor. CD27-binding protein (SIVA), a proapoptotic protein, can bind to this receptor and is thought to play an important role in the apoptosis induced by this receptor.

Species/Host: HEK293

Molecular Weight: The protein has a predicted molecular mass of 65-72 kDa after removal of the signal peptide.

Molecular Characterization: CD27(Ala20-Arg191) mFc(Pro99-Lys330) 6xHis

Purity: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% - 8% trehalose is added as protectants before lyophilization.

Storage & Shipping: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.




Figure 1. Human CD27 Protein, mFc-His Tag on SDS-PAGE under reducing condition.

Human CD27, mFc-His Tagged protein ELISA

0.2 μ g of CD27, mFc-His Tagged protein per well




Figure 2. ELISA plate pre-coated by 2 μ g/ml (100 μ l/well) Human CD27, mFc-His tagged protein (11141) can bind Anti-CD27 Neutralizing antibody 28027 in a linear range of 0.64-16.0 ng/ml.




Figure 3. ELISA plate pre-coated by 2 μ g/ml (100 μ l/well) Human CD27, mFc-His tagged protein (11141) can bind Human CD70, hFc-His tagged protein 11294 in a linear range of 3.2-80 ng/ml.




Figure 4. Flow cytometry analysis with 15 μ g/ml Human CD27 Protein, mFc-His tag (11141) on Expi293 cells transfected with human CD70 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).