

## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010

## HUMAN CD47 PROTEIN, MFC-HIS TAG

Cat.#: 11136 Product Name: Human CD47 Protein Size: 10 μg, 50 μg and 100 μg Synonyms: CD47;MER6;IAP;OA3 Target: CD47 UNIPROT ID: Q08722

Description: Recombinant human CD47 protein with C-terminal mouse Fc and 6xHis tag

**Background:** Leukocyte surface antigen CD47 is also known as Antigenic surface determinant protein OA3, Integrinassociated protein (IAP) and Protein MER6. CD47 contains 1 Ig-like V-type (immunoglobulin-like) domain. CD47 is very broadly distributed on normal adult tissues. CD47 has a role in both cell adhesion by acting as an adhesion receptor for THBSI on platelets, and in the modulation of integrins and plays an important role in memory formation and synaptic plasticity in the hippocampus by similarity. CD47 is the receptor for SIRPA, binding to which prevents maturation of immature dendritic cells and inhibits cytokine production by mature dendritic cells. CD47 Interaction with SIRPG mediates cell-cell adhesion, enhances superantigen-dependent T-cell-mediated proliferation and costimulates T-cell activation. **Species/Host:** HEK293

**Molecular Weight:** The protein has a predicted molecular mass of 60 kDa after removal of the signal peptide. **Molecular Characterization:** CD47(Gln19-Gln141) CD47(Val198-Ala207) CD47(Ser257-Gly268) mFc(Pro99-Lys330) 6×His tag

**Purity:** The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining. **Formulation & Reconstitution:** Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

**Storage & Shipping:** Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.



## **Product Description**

Pioneering GTPase and Oncogene Product Development since 2010




Figure 1. Human CD47, mFc-His Tag on SDS-PAGE under reducing condition.




Figure 2. ELISA plate pre-coated by 2 µg/ml (100 µl/well) Human CD47, mFc-His tagged protein (11136) can bind its native ligand Human SIRPI±, hFc-His tagged protein 11137 in a linear range of 3.3-26.37 ng/ml.




Figure 3. ELISA plate pre-coated by 2 µg/ml (100 µl/well) Human CD47, mFc-His tagged protein (11136) can bind Anti-CD47 (Neutralizing antibody clone Hu5F9-G4) 28010 in a linear range of 0.061-1.606 ng/ml.




Figure 4. ELISA plate pre-coated by 1 µg/ml (100 µl/well) Human CD47, mFc-His tagged protein (11136) can bind Anti-CD47 Neutralizing antibody 28058 in a linear range of 3.2-80.0 ng/ml.