

## HUMAN CD70 PROTEIN, HFC-HIS TAG

**Cat.#:** 11294

**Product Name:** Human CD70 Protein

**Size:** 10 µg, 50 µg and 100 µg

**Synonyms:** CD70;D27LG;NFSF7;NFSF7G;D27L

**Target:** CD70

**UNIPROT ID:** P32970

**Description:** Recombinant human CD70 protein with C-terminal human Fc and 6xHis tag

**Background:** The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T and B lymphocytes. It induces proliferation of costimulated T cells, enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin synthesis.

**Species/Host:** HEK293

**Molecular Weight:** The protein has a predicted molecular mass of 44.1 kDa after removal of the signal peptide.

**Molecular Characterization:** CD70(Gln39-Pro193) hFc(Glu99-Ala330) 6xHis

**Purity:** The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

**Formulation & Reconstitution:** Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

**Storage & Shipping:** Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.




Figure 1. Human CD70 Protein, hFc-His Tag on SDS-PAGE under reducing condition.




Figure 3. Flow cytometry analysis with 15  $\mu\text{g}/\text{ml}$  Human CD70 Protein, hFc-His tag (11294) on Expi293 cells transfected with human CD27 (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).