

Product Description

Pioneering GTPase and Oncogene Product Development since 2010

HUMAN FGF-9 PROTEIN

Cat.#: 12106

Product Name: Human FGF-9 Protein **Size:** 10 μg, 50 μg and 100 μg

Synonyms: Fibroblast Growth Factor 9;FGF-9;Glia-Activating Factor;GAF;Heparin-Binding Growth Factor 9;HBGF-9;FGF9

Target: FGF-9
UNIPROT ID: P31371

Description: Recombinant Human Fibroblast Growth Factor 9 is produced by our E.coli expression system and the target

gene encoding Met1-Ser208 is expressed.

Background: Fibroblast Growth Factor 9 (FGF-9) belongs to the Fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. FGF-9 plays an important role in the regulation of embryonic development, cell proliferation, cell differentiation and cell migration. In addition, FGF-9 may have a role in glial cell growth and differentiation during development, gliosis during repair and regeneration of brain tissue after damage, differentiation and survival of neuronal cells, and growth stimulation of glial tumors.

Species/Host: E.coli

Molecular Weight: 23.44 KDa

Molecular Characterization: Not available

Purity: Greater than 95% as determined by reducing SDS-PAGE.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

Storage & Shipping: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.




Figure 1. Greater than 95% as determined by reducing SDS-PAGE.




Figure 2. Measured in a cell proliferation assay using Balb/3T3 mouse embryonic fibroblast cells. The ED50 for this effect is 1-5 ng/ml.