

HUMAN GDNF PROTEIN, HFC TAG

Cat.#: 11645

Product Name: Human GDNF Protein

Size: 10 µg, 50 µg and 100 µg

Synonyms: ATF;ATF1;ATF2;HFBI-GDNF;HSCR3

Target: GDNF

UNIPROT ID: P39905

Description: Recombinant Human GDNF with C-terminal human Fc tag

Background: This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate each subunit of the disulfide-linked homodimer. The recombinant form of this protein, a highly conserved neurotrophic factor, was shown to promote the survival and differentiation of dopaminergic neurons in culture, and was able to prevent apoptosis of motor neurons induced by axotomy. This protein is a ligand for the product of the RET (rearranged during transfection) protooncogene. Mutations in this gene may be associated with Hirschsprung disease and Tourette syndrome. This gene encodes multiple protein isoforms that may undergo similar proteolytic processing. [provided by RefSeq, Aug 2016]

Species/Host: HEK293

Molecular Weight: The protein has a predicted molecular mass of 41.2 kDa after removal of the signal peptide. The apparent molecular mass of GDNF-hFc is approximately 55-70 kDa due to glycosylation.

Molecular Characterization: GDNF(Ser78-Ile211) hFc(Glu99-Ala330)

Purity: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

Storage & Shipping: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.




Figure 1. Human FOLR2 Protein, hFc Tag on SDS-PAGE under reducing condition.