

HUMAN PILRA PROTEIN, HFC TAG**Cat.#:** 11931**Product Name:** Human PILRA Protein**Size:** 10 µg, 50 µg and 100 µg**Synonyms:** FDF03**Target:** PILRA**UNIPROT ID:** Q9UKJ1**Description:** Recombinant Human PILRA Protein with C-terminal human Fc tag

Background: Cell signaling pathways rely on a dynamic interaction between activating and inhibiting processes. SHP-1-mediated dephosphorylation of protein tyrosine residues is central to the regulation of several cell signaling pathways. Two types of inhibitory receptor superfamily members are immunoreceptor tyrosine-based inhibitory motif (ITIM)-bearing receptors and their non-ITIM-bearing, activating counterparts. Control of cell signaling via SHP-1 is thought to occur through a balance between PILRA α -mediated inhibition and PILRA β -mediated activation. These paired immunoglobulin-like receptor genes are located in a tandem head-to-tail orientation on chromosome 7. This particular gene encodes the ITIM-bearing member of the receptor pair, which functions in the inhibitory role. Alternative splicing has been observed at this locus and three variants, each encoding a distinct isoform, are described. [provided by RefSeq, Jul 2008]

Species/Host: HEK293

Molecular Weight: The protein has a predicted molecular mass of 46.2 kDa after removal of the signal peptide. The apparent molecular mass of PILRA-hFc is approximately 55-70 kDa due to glycosylation.

Molecular Characterization: PILRA(Gln20-Glu195) hFc(Glu99-Ala330)

Purity: The purity of the protein is greater than 95% as determined by SDS-PAGE and Coomassie blue staining.

Formulation & Reconstitution: Lyophilized from nanodisc solubilization buffer (20 mM Tris-HCl, 150 mM NaCl, pH 8.0). Normally 5% – 8% trehalose is added as protectants before lyophilization.

Storage & Shipping: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing). Lyophilized proteins are shipped at ambient temperature.




Figure 1. Human PILRA Protein, hFc Tag on SDS-PAGE under reducing condition.