

## IDH2(R172H)

### IDH2(R172H)

**Cat. #::** 26084

**Gene Symbol:** IDH2

**Description:** Anti-IDH2(R172H) Mouse Monoclonal Antibody

**Background:** Isocitrate dehydrogenase (IDH) catalyzes the oxidative decarboxylation of isocitrate to 2-oxoglutarate. Mutations of Arg-172 in IDH2 to Cys, His, Leu or Ser abolish magnesium binding and the conversion of isocitrate to alpha-ketoglutarate. Instead, alpha-ketoglutarate is converted to R-2-hydroxyglutarate. Elevated levels of R-2-hydroxyglutarate are correlated with an elevated risk of malignant brain tumors.

**Immunogen:** A synthetic peptide from the internal region of human IDH2 which includes the mutation of R172H.

**Applications:** ELISA, WB, IHC, IF

**Recommended Dilutions:**

ELISA: 1:1000-1:5000

WB: 1:100-1:1000

IHC: 1:50-1:100

IF: 1:50-1:100

**Concentration:** 1 mg/ml

**Host Species:** Mouse

**Format:** Liquid

**Clonality:** Monoclonal

**Isotype:** IgG

**Purity:** Purified from ascites


**Preservative:** No

**Constituents:** PBS (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150 mM NaCl, 50% glycerol

**Species Reactivity:** Anti-IDH2(R172H) antibody recognizes IDH2(R172H) of vertebrates.

**Storage Conditions:** Store at -20°C. Avoid repeated freezing and thawing.


### Western blot:



## Western blot analysis of recombinant IDH2(R172H) and wild type IDH2 proteins.

Purified his-tagged IDH2(R172H) protein (Cat. # 10201) and wild type IDH2 protein (Cat. # 10200) were blotted with anti-IDH2(R172H) monoclonal antibody (Cat. # 26084) (left panel). Proteins loading were blotted with anti-IDH2 monoclonal antibody (right panel).

## Immunofluorescence:



## Immunofluorescence of cells expressing IDH2 proteins with Anti-IDH2(R172H) antibody.

HEK293T cells were transfected with pCDNA3-GFP-IDH2 (WT) plasmid (left column) or pCDNA3-GFP-IDH2(R172H) plasmid (right column), then fixed and stained with Anti-IDH2(R172H) monoclonal antibody (Cat. #26084).