

IDH2(R172H)**IDH2(R172H)****Cat. #::** 26084**Gene Symbol:** IDH2**Description:** Anti-IDH2(R172H) Mouse Monoclonal Antibody**Background:** Isocitrate dehydrogenase (IDH) catalyzes the oxidative decarboxylation of isocitrate to 2-oxoglutarate. Mutations of Arg-172 in IDH2 to Cys, His, Leu or Ser abolish magnesium binding and the conversion of isocitrate to alpha-ketoglutarate. Instead, alpha-ketoglutarate is converted to R-2-hydroxyglutarate. Elevated levels of R-2-hydroxyglutarate are correlated with an elevated risk of malignant brain tumors.**Immunogen:** A synthetic peptide from the internal region of human IDH2 which includes the mutation of R172H.**Applications:** ELISA, WB, IHC, IF**Recommended Dilutions:**


ELISA: 1:1000–1:5000

WB: 1:100–1:1000

IHC: 1:50–1:100

IF: 1:50–1:100


Concentration: 1 mg/ml**Host Species:** Mouse**Format:** Liquid**Clonality:** Monoclonal**Isotype:** IgG**Purity:** Purified from ascites**Preservative:** No**Constituents:** PBS (without Mg^{2+} and Ca^{2+}), pH 7.4, 150 mM NaCl, 50% glycerol**Species Reactivity:** Anti-IDH2(R172H) antibody recognizes IDH2(R172H) of vertebrates.**Storage Conditions:** Store at $-20^{\circ}C$. Avoid repeated freezing and thawing.**Western blot:**



Western blot analysis of recombinant IDH2(R172H) and wild type IDH2 proteins.

Purified his-tagged IDH2(R172H) protein (Cat. # 10201) and wild type IDH2 protein (Cat. # 10200) were blotted with anti-IDH2(R172H) monoclonal antibody (Cat. # 26084) (left panel). Proteins loading were blotted with anti-IDH2 monoclonal antibody (right panel).

Immunofluorescence:



Immunofluorescence of cells expressing IDH2 proteins with Anti-IDH2(R172H) antibody.

HEK293T cells were transfected with pCDNA3-GFP-IDH2 (WT) plasmid (left column) or pCDNA3-GFP-IDH2(R172H) plasmid (right column), then fixed and stained with Anti-IDH2(R172H) monoclonal antibody (Cat. #26084).