

IL5RA (DMC393) IGG1 CHIMERIC MAB**Cat.#:** 28211**Product Name:** Anti-IL5RA(DMC393) IgG1 Chimeric Monoclonal Antibody**Synonyms:** IL5Ra;CD125;IL-5 R alpha**Description:** Anti-IL5RA antibody(DMC393) IgG1 Chimeric Monoclonal Antibody

Background: The protein encoded by this gene is an interleukin 5 specific subunit of a heterodimeric cytokine receptor. The receptor is comprised of a ligand specific alpha subunit and a signal transducing beta subunit shared by the receptors for interleukin 3 (IL3); colony stimulating factor 2 (CSF2:GM-CSF); and interleukin 5 (IL5). The binding of this protein to IL5 depends on the beta subunit. The beta subunit is activated by the ligand binding; and is required for the biological activities of IL5. This protein has been found to interact with syndecan binding protein (syntenin); which is required for IL5 mediated activation of the transcription factor SOX4. Several alternatively spliced transcript variants encoding four distinct isoforms have been reported.

Applications: Flow Cyt**Recommended Dilutions:** Flow Cyt 1:100**Host Species:** Rabbit**Isotype:** Rabbit:Human Fc chimeric IgG1**Purification:** Purified from cell culture supernatant by affinity chromatography**Species Reactivity:** Human IL5RA**Constituents:** Lyophilized from sterile PBS, pH 7.4. 5 % – 8% trehalose is added as protectants before lyophilization.**Storage & Shipping:** Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).




Figure 1. Flow cytometry analysis with Anti-IL5RA (DMC393) on Expi293 cells transfected with human IL5RA (Blue histogram) or Expi293 transfected with irrelevant protein (Red histogram).